

Table 1. Average nightly number of brown marmorated stink bug (BMSB) captured in eastern Virginia black light traps, 2012 season.

Nightly BMSB trap catch average for week ending:											
Location (county-town)	5/31	6/07	6/14	6/21	6/28	7/05	7/12	7/19	7/26	8/02	8/09
Southeast											
Accomack-Painter.....	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Chesapeake-Ballahack Rd.	n/a	0.0	0.0	0.0	0.0	n/a	0.0	0.0	n/a	n/a	n/a
Dinwiddie-Old Hickory	0.0	0.0	0.0	0.0	n/a	0.8	1.9	1.9	n/a	n/a	n/a
Dinwiddie-North.....	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Isle of Wight	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.0	*
Petersburg	2.6	0.0	0.3	0.3	0.9	0.9	2.0	19.6	25.7	31.6	13.9
Prince George-Templeton.....	n/a	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0
Prince George-Disputanta.....	n/a	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9
Southampton	n/a	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Suffolk-Holland	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Surry	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.0
Sussex-Waverly	n/a	n/a	n/a	n/a	n/a	n/a	0.0	0.4	n/a	0.0	n/a
Virginia Beach-HRAREC.....	0.0	0.0	0.0	0.0	1.0	0.0	1.0	2.4	2.6	2.7	n/a
Virginia Beach-West Neck	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.0	n/a	0.0	n/a
North of James River											
Essex	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.0	0.0	0.0	n/a
Gloucester	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
King William	n/a	n/a	n/a	0.9	0.0	0.1	1.9	2.5	7.3	n/a	n/a
Northumberland	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
Middlesex	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
New Kent	n/a	n/a	n/a	n/a	n/a	n/a	0.0	4.3	n/a	0.6	n/a
Richmond Co.-Warsaw.	n/a	n/a	n/a	0.3	0.1	0.3	0.4	2.0	2.7	2.2	2.8

n/a = report not available.

*Isle of Wight had 1 suspected BMSB captured (awaiting species confirmation).