

Table 1. Average nightly number of *Helicoverpa zea* (corn earworm/bollworm) moths captured in eastern Virginia black light traps, 2017 season.

Nightly <i>H. zea</i> trap catch average for the week ending:																	
Location (county-town)	6/29	7/06	7/13	7/20	7/27	8/03	8/10	8/17	8/24	8/31	9/07	9/14	9/21	9/28	10/05	10/12	10/19
Chesapeake.....	n/a	n/a	0.0	2.0	n/a												
Dinwiddie-Old Hickory	n/a	n/a	3.6	5.2	10.4												
Dinwiddie-Ford.....	n/a	n/a	n/a	n/a	n/a												
Hanover.....	n/a	0.2	2.7	1.1	20.4												
Prince George-Templeton	n/a	n/a	n/a	0.8	2.3												
Prince George-Disputanta	n/a	n/a	n/a	0.3	4.6												
Richmond Co.-Warsaw .	1.3	0.0	0.0	0.4	7.1												
Southampton-Courtland	n/a	0.6	0.0	0.0	2.1												
Suffolk-Holland.....	n/a	n/a	n/a	8.5	65.3												

n/a = report not available.

Table 2. Average nightly number of brown marmorated stink bugs (BMSB) captured in eastern Virginia black light traps, 2017 season.

Nightly BMSB trap catch average for the week ending:																	
Location (county-town)	6/29	7/06	7/13	7/20	7/27	8/03	8/10	8/17	8/24	8/31	9/07	9/14	9/21	9/28	10/05	10/12	10/19
Chesapeake.....	n/a	n/a	0.0	0.0	n/a												
Dinwiddie-Old Hickory	n/a	n/a	0.0	0.0	0.0												
Dinwiddie-Ford.....	n/a	n/a	n/a	n/a	n/a												
Hanover.....	n/a	1.0	1.9	2.7	10.0												
Prince George-Templeton	n/a	n/a	n/a	0.2	0.0												
Prince George-Disputanta	n/a	n/a	n/a	0.0	0.7												
Richmond Co.-Warsaw .	0.0	0.0	0.1	1.4	1.9												
Southampton-Courtland	n/a	0.0	0.0	0.0	0.0												
Suffolk-Holland.....	n/a	n/a	n/a	0.0	0.1												

n/a = report not available.